

---

Citation:

Price, OJ and Tsakirides, C and Gray, M and Stavropoulos-Kalinoglou, A (2018) ACSM pre-participation health screening guidelines: a UK university cohort perspective. *Medicine and Science in Sports and Exercise*. ISSN 1530-0315 DOI: <https://doi.org/10.1249/MSS.0000000000001868>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/5566/>

Document Version:

Article (Accepted Version)

---

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

**ACSM PRE-PARTICIPATION HEALTH SCREENING GUIDELINES: A UK  
UNIVERSITY COHORT PERSPECTIVE**

Oliver J. Price PhD\*, Costas Tsakirides MSc, Mike Gray MSc, Antonios Stavropoulos-  
Kalinoglou PhD\*

Carnegie School of Sport, Leeds Beckett University, Leeds, United Kingdom (UK).

**\*Authors contributed equally**

**Corresponding author:**

Dr Oliver J. Price BSc (Hons.) MRes PhD FHEA

Carnegie School of Sport, Leeds Beckett University,

Leeds, Headingley Campus, LS6 3QT, UK

Tel: +44 (0)113 8123 532

Email: [o.price@leedsbeckett.ac.uk](mailto:o.price@leedsbeckett.ac.uk)

\*No funding was received for this work (from NIH, Wellcome Trust, HHMI, or any others).

\*The authors have no real or perceived conflict of interest in respect of this manuscript.

**Manuscript word count:** 2954; **Abstract word count:** 267.

**Running title:** Exercise preparticipation health screening.

## ABSTRACT

**Purpose:** Pre-participation health screening is recommended to detect individuals susceptible to serious adverse cardiovascular complications during exercise. Although expert opinion and best available scientific evidence have informed recent modifications, there remains limited experimental data to support or refute current practice. We therefore aimed to quantify the impact of change to the ACSM pre-participation health screening guidelines on risk classification and referral for medical clearance in a large cohort of undergraduate university students. **Methods:** Participants attended the laboratory on a single occasion to undergo pre-participation health screening. Information concerning health status was obtained via self-report questionnaire and objective physiological assessment with all data recorded electronically and evaluated against ACSM screening guidelines (9<sup>th</sup> and 10<sup>th</sup> Edition). **Results:** Five-hundred and fifty-three students completed the study. The 9<sup>th</sup> Edition screening guidance resulted in eighty-two (15%) subjects classified as high-risk, almost one quarter (24%) classified as moderate-risk, and almost two-thirds (61%) classified as low-risk. In comparison, the updated 10<sup>th</sup> Edition screening guidance resulted in a significant reduction in those previously classified as either high-risk (5%) or moderate risk (2%), respectively. The majority of subjects (93%) were therefore cleared to begin a structured exercise programme. Taken together, approximately one-third (32%) fewer medical referrals were required when applying the updated 10<sup>th</sup> Edition guidance ( $\chi^2(4) = 247.7, P < 0.001$ ). **Conclusion:** The updated ACSM 10<sup>th</sup> Edition pre-participation screening guidance reduces medical referrals by approximately one-third. These findings are in keeping with previous reports and thus serve to consolidate and justify recent modification - particularly when applied to young adult or adolescent populations. The findings and arguments presented should be used to refine and inform future guidance.

**Key words:** ACSM, Exercise, Health, Physical Activity, Pre-participation, Screening.

## 1 INTRODUCTION

2 Regular physical activity and structured exercise training are widely accepted to promote health  
3 and well-being in addition to lowering the risk of developing chronic disease such as  
4 cardiovascular disease (CVD), diabetes mellitus and cancers across the lifespan (1). However,  
5 despite recognition of health benefit (i.e. reduced morbidity and mortality) (2), a large  
6 proportion (~30%) of the global population remain sedentary (3, 4). On this basis, and to reduce  
7 healthcare costs attributed to sedentary living (5), the World Health Organisation (WHO) have  
8 recently targeted a reduction in physical inactivity over the next decade as a key global health  
9 priority (6-9).

10 The American College of Sports Medicine (ACSM) currently advise completing at least 30  
11 minutes of moderate intensity activity 5 days per week, or 20 minutes of vigorous intensity  
12 activity 3 days per week to optimise and maintain health (10). However, prior to initiating an  
13 exercise programme, irrespective of age, it is recommended that pre-participation health  
14 screening is conducted to detect individuals susceptible to serious adverse cardiovascular  
15 complications; i.e. sudden cardiac death and/or acute myocardial infarction during or post  
16 exercise (10).

17 Historically, pre-participation health screening, as described in the 9<sup>th</sup> Edition and previous  
18 versions of the ACSM Guidelines to Exercise Testing and Prescription focused on: (a) risk  
19 stratification calculated according to CVD risk factors and (b) the presence of signs or  
20 symptoms of and/or history of cardiovascular, pulmonary, renal, liver or metabolic disease.  
21 This approach permitted recommendations concerning suitability to commence exercise or the  
22 requirement for further clinical assessment (i.e. referral to a physician to obtain medical  
23 clearance) (11).

More recently, however, it has been acknowledged that employing this approach may contribute to excessive medical referrals thereby placing unnecessary burden on healthcare infrastructure, and equally, may create a barrier to physical activity and exercise engagement (12). In an attempt to simplify guidance, and address these concerns, a panel of experts convened by the ACSM published modifications to the pre-participation health screening criteria in 2015 (subsequently incorporated in the 10<sup>th</sup> Edition of the Guidelines for Exercise Testing and Prescription) (10). Accordingly, screening guidance currently centres on: (a) current level of physical activity, (b) the presence of known cardiovascular, metabolic, or renal disease (or signs or symptoms of these diseases), and (c) the desired intensity of the exercise bout / programme (10).

Although expert opinion and best available scientific evidence have informed these modifications, there remains limited experimental data to support or refute current practice, particularly when applied to young adult (i.e. <40 years) or adolescent populations (13). This is important given pre-participation health screening is recommended within universities and colleges (notably across exercise science courses and sports teams), and as part of inductions to gymnasia and fitness centres around the globe.

The primary aim of this study was therefore to quantify the impact and examine the implications of change to the ACSM pre-participation health screening guidelines on risk classification and referral for medical clearance in a large cohort of UK-based undergraduate university students.

A secondary aim was to evaluate the requirement for widespread screening in this context by characterising the prevalence of CVD risk and signs/symptoms or history of non-communicable disease. We hypothesised that the updated 10<sup>th</sup> Edition guidance would reduce the percentage of medical referrals whilst providing a cost-effective and time-efficient approach to pre-participation health screening in young adult populations.

## **METHODS**

### **Study population and experimental design**

Six-hundred and five undergraduate university sport and exercise science students were enrolled into the study between 2015-17. In accordance with the Leeds Beckett University (UK) local pre-exercise risk assessment and risk management policy, all participants attended the laboratory on a single occasion to undergo pre-participation health screening. Information concerning health status was obtained via self-report questionnaire and objective physiological assessment with all data recorded electronically. In accordance with ACSM 9<sup>th</sup> Edition guidance (described below) - a finger-prick capillary blood sample was obtained for the assessment of blood glucose and total cholesterol for subjects who presented with  $\geq 1$  CVD risk factor. All testing was conducted by trained personnel and evaluated against ACSM pre-participation screening guidelines (9<sup>th</sup> and 10<sup>th</sup> Edition). Data were analysed retrospectively by two independent reviewers (OP and AS) to complete risk classification and determine screening outcome. All tests and procedures were approved by the local research ethics committee (IRB ethics ID: 50858) and all subjects provided written informed consent (Figure 1).

### **Pre-participation health screening guidance**

#### ***9<sup>th</sup> Edition***

The ACSM 9<sup>th</sup> Edition algorithm classifies individuals according to risk; i.e. low, moderate, and high. Screening outcome is determined based on the sum of CVD risk factors; i.e. age, family history of heart disease, smoking status, physical activity, obesity (based on BMI and/or waist circumference), dyslipidaemia, dysglycaemia and resting blood pressure values, and presence or absence of signs or symptoms, or known cardiovascular, *pulmonary*, renal, liver or metabolic disease with recommendations for medical clearance based on the desired intensity of exercise (Figure 2A).

## **10th Edition**

The ACSM 10th Edition algorithm classifies individuals as physically active or inactive according to the level of exercise or physical activity engagement over the past 3 months (defined as planned, structured physical activity of at least 30 min duration at moderate intensity on at least 3 days per week). Screening outcome is determined based on presence or absence of known cardiovascular, metabolic, or renal disease and presence or absence of signs or symptoms of these conditions with recommendations for medical clearance based on the desired intensity of exercise (Figure 2B).

### **Risk stratification and screening outcome (9th vs. 10th Edition)**

To allow a direct comparison of guidance, risk stratification and screening outcome was classified as (a) low-risk (colour code: green) (i.e. medical clearance not necessary), (b) moderate-risk (colour code: yellow) (i.e. medical clearance only required for vigorous exercise), (c) high-risk (colour code: red) (i.e. medical clearance recommended prior to any physical activity) (Figure 2).

### **Statistical analysis**

Referral percentages were calculated and stratified with cross-tabulation conducted to compare screening outcome. Independent t-tests (continuous variables) and Chi-squared tests (categorical variables) were employed to evaluate between group (sex) differences. All other data were reported descriptively according to prevalence of CVD risk factors, signs and symptoms of disease and prior history of non-communicable disease. Data were analysed using SPSS Statistics statistical software package Version 24 (SPSS Inc., Chicago, IL).  $P < 0.05$  was considered statistically significant. Data are reported as mean  $\pm$  SD unless otherwise stated.

## RESULTS

### Study population

Of the 605 subjects recruited to this study, 553 (male  $n = 393$ ) completed the study, while 52 (male  $n = 43$ ) were excluded from the analysis due to incomplete datasets. Subjects' clinical characteristics are presented in Table 1.

### Comparison of risk stratification and screening outcome

The 9<sup>th</sup> Edition screening guidance resulted in eighty-two (15%) subjects classified as high-risk, one-hundred and thirty five (24%) classified as moderate-risk, and three hundred thirty three (61%) classified as low-risk. In comparison, the updated 10<sup>th</sup> Edition screening guidance resulted in twenty-seven classified as high-risk (5%) and thirteen classified as moderate risk (2%), respectively. All subjects originally classified as moderate risk according to 9<sup>th</sup> Edition guidance were re-classified as low risk when applying the 10<sup>th</sup> Edition guidance. The remaining five hundred and thirteen (93%) were therefore cleared to begin a structured exercise programme in accordance with the 10<sup>th</sup> edition of the ACSM's Guidelines for Exercise Testing and Prescription. Taken together, almost one-third (32%) fewer medical referrals were required when applying the updated 10<sup>th</sup> Edition guidance ( $\chi^2(4) = 247.7$ ,  $P < 0.001$ ) (Figure 3A).

### High-risk re-classification

Of the eighty-two subjects (15% of all participants) originally classified as high-risk, twenty-seven (5%) remained the same and required referral for medical clearance due to a sign and/or symptom of cardiovascular, metabolic or renal disease. Thirteen subjects (16%) were re-classified as moderate risk due to being physically active and having known cardiovascular, metabolic or renal disease but also being asymptomatic. Forty-two subjects (51%) were re-classified as low-risk due to guidance modification concerning pulmonary and liver (Figure 3B). Specifically, thirty-eight subjects (46%) diagnosed with asthma, one subject (1%)



diagnosed with cystic fibrosis, one subject (1%) diagnosed with interstitial lung disease, and two subjects (2%) diagnosed with liver disease were reclassified as low-risk (Table 2).

### **CVD risk factors, signs and/or symptoms and history of disease**

Prevalence of family history of CVD, physical inactivity, obesity (based on BMI and waist circumference thresholds), dyslipidaemia and dysglycaemia were not different between sexes ( $P>0.05$ ). In contrast, smoking status ( $\chi^2(1) = 9.172, p=0.002$ ) and evidence of hypertension ( $\chi^2(1) = 5.995, p=0.014$ ) were higher in males. Descriptive statistics and prevalence data for CVD are provided in Table 3. Signs and/or symptoms of disease were not different between sexes ( $P>0.05$ ). For the majority of diseases, no difference was observed between sexes ( $P>0.05$ ). Descriptive statistics and prevalence data for signs and/or symptoms and history of non-communicable disease are provided in Table 2.

### **Blood sample analysis ( $n = 135$ )**

One hundred and thirty-five subjects (male:  $n = 102$ ) (24%) presented with  $\geq 1$  CVD risk factor and therefore provided a capillary blood sample in accordance with 9<sup>th</sup> Edition screening guidance. Of these, only five subjects (4%) presented with dyslipidaemia and only two subjects (1%) presented with dysglycaemia. In contrast, the other one hundred and twenty-eight subjects (95%) (1 CVD risk factor) were confirmed as low-risk following determination of normal resting blood lipid and glucose profiles (Table 3).

## DISCUSSION

In support of our hypothesis, the present study provides the first experimental evidence to confirm that when employing the revised ACSM 10<sup>th</sup> Edition pre-participation screening guidance, almost one-third (32%) fewer university students were referred for medical clearance. The implications of our findings can be considered two-fold: firstly, reducing the impact of unnecessary or excessive medical referrals on primary healthcare infrastructure (i.e. cost, time and resource), and secondly, reducing barriers to exercise and physical activity engagement. The latter is of particular importance given the strong association between exercise engagement in early life and physical activity status across the lifespan (14).

The majority of research to date investigating pre-participation health screening has primarily evaluated individuals considered to be at 'heightened' risk of cardiovascular complications during exercise (i.e. mid-adult or elderly age-groups) (13, 15). For example, a recent retrospective analysis by Whitfield and colleagues compared referral patterns between the ACSM 10<sup>th</sup> Edition guidance with previously validated (and widely implemented) pre-participation questionnaires (i.e. AAPQ and PAR-Q) in a large representative sample ( $n = 6785$ ) of United States adults (13). Similar to the present study, it was found that the 10<sup>th</sup> Edition guidance resulted in significantly fewer (~38%) medical referrals. The sample size of this comparative analysis is clearly a study strength, however it is important to acknowledge that several symptoms contributing to the ACSM guidance were not assessed, with all adults aged 40-years or older (13).

In contrast to this approach, for the first time, our study provides a direct comparison of screening methods (i.e. ACSM 9<sup>th</sup> vs. 10<sup>th</sup> Edition) applicable to younger age-groups (i.e. young adult and adolescent populations). It is our view, that the data from the present study should therefore be interpreted to consolidate previous findings and interpreted to support

recent modification and inform potential revisions to future ACSM guidance. Taken together, the simplicity, cost-effectiveness and short time-frame to complete and obtain a screening outcome when applying the 10<sup>th</sup> Edition guidance provides strong justification to employ widespread screening (irrespective of age) prior to beginning an exercise programme.

The lower proportion of referrals and change to screening outcome in the present study is attributable to guidance modification concerning: (a) evaluating physical activity engagement, (b) removal of pulmonary and liver disease from screening outcome, and (c) discounting CVD risk factors (10). Of the eighty-two subjects that were originally classified as high-risk according to 9th Edition guidance, twenty-seven (15%) remained unchanged due to reporting a sign and/or symptom of cardiovascular, metabolic or renal disease. However, thirteen (16%) (presenting with known cardiovascular, metabolic or renal disease and asymptomatic) were re-classified to moderate risk due to physical activity either meeting or exceeding current recommendations (10). As the absolute and relative risks of acute cardiovascular events during exercise are known to be very low in asymptomatic individuals (16-18), re-classification and clearance to begin moderate intensity exercise (40-60% aerobic capacity) is certainly logical when applied to physically active individuals (irrespective of disease status).

It is important to note that self-report physical activity data acquired through consultation and questionnaire has previously been reported to provide poor validity and reliability when compared to device-based measures of human movement (e.g. pedometers and accelerometers etc.) (19). Indeed the most comprehensive appraisal of the literature to date concluded that self-report frequently results in both an over and under-estimation of physical activity (20). These findings therefore have important ramifications for risk assessment when employing 10<sup>th</sup> Edition guidance in the context of health screening and exercise prescription. Although the prominence of physical activity in the 10th Edition positively contributes to the observed

reduction in medical referrals, limitations associated with self-report should be considered during consultation. To optimise the accuracy of estimated daily physical activity, what constitutes ‘planned, structured physical activity’ should be clearly defined, and it is our view that supplementary or follow-up questions should be considered and encouraged moving forward.

The decision to no longer automatically refer those with pulmonary disease had a significant impact on screening outcome. As evidenced by our findings, of those previously classified as high-risk, almost half (49%) were re-classified as low-risk when discounting pulmonary disease (asthma: 95%, cystic fibrosis: 2.5% and interstitial lung disease: 2.5%). In support of this modification, it is worth noting that pulmonary disease is not associated with an increased risk of cardiovascular complications during exercise, rather, long-term physical inactivity, due to fear of provoking exertional breathlessness or exacerbation, is more likely to increase the relative-risk of an adverse cardiovascular event in this population (21). The overall prevalence of asthma (~7%) in our cohort is in keeping with current UK estimates (~8%) (22). It is important to acknowledge however that asthma is a complex disease consisting of several variants with unique underlying pathophysiology (23). In the context of pre-participation health screening, asthma sub-type and clinical control status should also be considered before granting clearance to commence vigorous or maximal intensity exercise. Caution is therefore advised when screening individuals predisposed to exercise-induced asthma, or those with severe, uncontrolled, or difficult to treat airways disease (24). Irrespective of this recommendation, the reduced referral pattern should help to deter physical inactivity in pulmonary disease patients. Indeed, the importance of regular exercise and physical activity in the context of pulmonary rehabilitation is well-established (25, 26).

Perhaps unsurprisingly liver disease was only found to have a minor impact on screening outcome in our young adult population. However, considering the growing global epidemic of obesity and associated metabolic disease (largely attributed to modifiable lifestyle factors such as diet, alcohol consumption, and physical inactivity), it is likely that a similar reduction in referral proportions related to liver disease would be observed when applied to older or ageing populations (27). Similarly, for the most part, the prevalence of CVD risk factors in our cohort was relatively low (<5%) with no difference observed between sexes.

The suggestion of excessive referrals in the past is supported by our data given the majority of our population (95%) who presented with  $\geq 1$  CVD risk factor (following the screening questionnaire and measurements of anthropometry and resting blood pressure) returned to low-risk following blood analysis (i.e. normal resting blood glucose and lipid profiles). Of concern, the incidence of smoking status (9%) and evidence of hypertension (14%) was high in males, which clearly has ramifications for long-term respiratory health (28) and the development of CVD and heightened relative risk of mortality in later life (29). Finally, the incidence of signs and/or symptoms or history of non-communicable disease was low when compared against current estimates from the global population (30).

### **Methodological considerations and future research**

Despite our encouraging findings, it is important to recognise and consider potential limitations to the updated 10<sup>th</sup> Edition guidance. For example, current policy stipulates clearance to engage in ‘vigorous intensity exercise’ defined as  $\geq 60\%$  aerobic capacity. Whilst evidence indicates that the absolute risk of cardiovascular complications during high-intensity exercise is low, the relative risk of sudden cardiac death and/or acute myocardial infarction during or post exercise increases (irrespective of age, current physical fitness or functional capacity) when approaching maximal exercise. To further mitigate risk; in circumstances where individuals

are required to exercise to volitional exhaustion (e.g. maximal oxygen uptake testing), it may be pragmatic to evaluate the accuracy of self-report information relevant to screening outcome (e.g. physical activity status) on a case-by-case basis before granting clearance. Although we recognise that ACSM pre-participation health screening guidelines are designed to detect individuals susceptible to serious adverse cardiovascular complications, it is also important to consider and rule-out other relevant conditions that may be made worse or increase the risk of injury during exercise. In support of this concept, musculoskeletal injury was reported in approximately 12% of our cohort which would have remained undetected without requesting further patient history. Finally, as with all screening criteria, a potential for false-positive or false-negative outcome exists when employing current guidance. To provide evidence-based recommendations and to justify benefit of widespread pre-participation health screening, validation against objective diagnostic work-up in those referred for medical clearance, and long-term follow-up of individuals cleared to exercise remains a research priority.

## **Conclusion**

In summary, in a large cohort of UK-based undergraduate university students, we have demonstrated that the ACSM 10th Edition guidance reduces medical referrals by approximately one-third. These findings are in-keeping with previous reports and thus serve to consolidate and justify recent modification - particularly when applied to young adult or adolescent populations. In order to continue to optimise the health, well-being and overall safety of individuals engaging in exercise, consideration of the findings and arguments presented should be used to refine and inform future guidance.

252 **TABLE FOOTNOTES**

253

254 **Table 1.** Subject clinical characteristics.

255 **Table 2.** Signs and symptoms and history of non-communicable disease (ACSM 9<sup>th</sup> and 10<sup>th</sup>  
256 Edition).

257 **Table 3.** Cardiovascular disease risk factors (ACSM 9<sup>th</sup> Edition)

258

259

260

## FIGURE LEGENDS

### Figure 1. Schematic detailing experimental design.

#### ACSM 9<sup>th</sup> Edition:

##### \*CVD risk factors

Age: men >45 years; women >55 years

Family history: myocardial infarction, coronary revascularisation, or sudden death before 55 year in father or other male first-degree relative or before 65 year in mother or other female first-degree relative

Cigarette smoking: Current cigarette smoker or those who quit within the previous 6-months or exposure to environmental tobacco smoke

Sedentary lifestyle: Not participating in at least 30 min of moderate intensity, physical activity (40% - <60% VO<sub>2</sub>R) on at least 3 day of the week for at least 3 months

Obesity: Body mass index  $\geq 30 \text{ kg.m}^2$  or waist circumference  $> 102 \text{ cm}$  for men and  $> 88 \text{ cm}$  for women

Hypertension: Systolic blood pressure  $\geq 140 \text{ mmHg}$  and/or diastolic  $\geq 90 \text{ mmHg}$ , confirmed by measurements on at least two separate occasions, or on anti-hypertensive medication

Dyslipidaemia: Low-density lipoprotein (LDL) cholesterol  $\geq 130 \text{ mg.dL}^{-1}$  ( $3.37 \text{ mmol.L}^{-1}$ ) or high-density lipoprotein (HDL) cholesterol  $< 40 \text{ mg.dL}^{-1}$  ( $1.04 \text{ mmol.L}^{-1}$ ) or on lipid-lowering medication. If total serum cholesterol is all that is available, use  $\geq 200 \text{ mg.dL}^{-1}$  ( $5.18 \text{ mmol.L}^{-1}$ )

Prediabetes: Impaired fasting glucose = fasting plasma glucose  $\geq 100 \text{ mg.dL}^{-1}$  ( $5.55 \text{ mmol.L}^{-1}$ ) and  $\leq 125 \text{ mg.dL}^{-1}$  ( $6.94 \text{ mmol.L}^{-1}$ ) or impaired glucose tolerance test = 2 hour values in oral glucose tolerance test  $\geq 140 \text{ mg.dL}^{-1}$  ( $7.77 \text{ mmol.L}^{-1}$ ) and  $\leq 199 \text{ mg.dL}^{-1}$  ( $11.04 \text{ mmol.L}^{-1}$ ) confirmed by measurements on at least two separate occasions.

Negative risk factors: High-density lipoprotein cholesterol  $\geq 60 \text{ mg.dL}^{-1}$  ( $1.55 \text{ mmol.L}^{-1}$ )

##### \*\*Signs or symptoms of disease

Pain, discomfort in the chest, neck, jaw, arms, or other areas that may result from ischemia

Shortness of breath at rest or with mild exertion

Dizziness or syncope

Orthopnoea or paroxysmal nocturnal dyspnoea

Ankle oedema

Palpitations or tachycardia

Intermittent claudication

Known heart murmur

Unusual fatigue or shortness of breath with usual activities

##### \*\*\*History of disease

Cardiovascular: Cardiac, peripheral vascular, or cerebrovascular disease

Pulmonary: Chronic obstructive pulmonary disease, asthma, interstitial lung disease or cystic fibrosis

Metabolic: Diabetes mellitus (types 1 and 2) or renal disease

#### ACSM 10<sup>th</sup> Edition:

##### †Participates in regular exercise

Performing planned, structured physical activity at least 30 min at moderate intensity on at least 3 days per week for at least the last 3 months

##### ††Signs or symptoms of disease

At rest or during activity; includes pain, discomfort in the chest, neck, jaw, arms, or other areas that may result from ischemia; shortness of breath at rest or with mild exertion; dizziness or syncope; orthopnoea or paroxysmal nocturnal dyspnoea; ankle oedema; palpitations or tachycardia; intermittent claudication; known heart murmur; or unusual fatigue or shortness of breath with usual activities

##### †††History of disease

Cardiac, peripheral vascular, or cerebrovascular disease

Type 1 and 2 diabetes mellitus



**Figure 2.** 9<sup>th</sup> (A) and 10<sup>th</sup> Edition (B) ACSM pre-participation health screening algorithm(s).

**Medical clearance:** Approval from a healthcare professional to engage in exercise

**Mod ex:** Moderate intensity exercise; 40% - <60% VO<sub>2</sub>R; 3 - <6 METS “An intensity that causes noticeable increases in heart rate and breathing”

**Vig ex:** Vigorous intensity exercise; ≥60% VO<sub>2</sub>R; ≥6 METS. “An intensity that causes substantial increases in heart rate and breathing”

**Figure 3A.** Screening outcome according to 9<sup>th</sup> and 10<sup>th</sup> Edition ACSM pre-participation health screening guidance.

**ACSM 9<sup>th</sup> Edition:** red - medical clearance recommended (*n* = 82; 15%); yellow - medical clearance required for vigorous intensity exercise only (*n* = 135; 24%); green - medical clearance not required (*n* = 333; 61%).

**ACSM 10<sup>th</sup> Edition:** red - medical clearance recommended (*n* = 27; 5%); yellow - medical clearance required for vigorous intensity exercise only (*n* = 13; 2%); green - medical clearance not required (*n* = 513; 93%)

**Figure 3B.** Re-classification of high-risk individuals (*n* = 82) according to updated pre-participation health screening guidelines (ACSM 10<sup>th</sup> Edition).

**ACSM 10<sup>th</sup> Edition:** red - medical clearance recommended (*n* = 27; 33%); yellow - medical clearance required for vigorous intensity exercise only (*n* = 13; 16%); green - medical clearance not required (*n* = 42; 51%).

**ACKNOWLEDGEMENTS**

The authors would like to acknowledge all Leeds Beckett University exercise physiologists and laboratory technicians who contributed to the acquisition of data.

**FUNDING STATEMENT**

Nil relevant.

**COMPETING INTERESTS**

The authors have no real or perceived conflict of interest in respect of this manuscript.

**CONTRIBUTION STATEMENT**

All authors were involved in the conception, design of the study, acquisition of data, drafting and critical revision of manuscript and final approval of the version to be published. The results of the study are presented clearly, honestly, and without fabrication or falsification.

**GUARANTOR STATEMENT**

OP and AS confirm full responsibility for the content of the manuscript, including data and analysis.

The results of the present study do not constitute endorsement by ACSM.

## REFERENCES

1. Garber CE, Blissmer B, Deschenes MR et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;43(7):1334-59.
2. Naci H, Ioannidis JP. Comparative effectiveness of exercise and drug interventions on mortality outcomes: metaepidemiological study. *BMJ.* 2013;347:f5577.
3. Hallal PC, Andersen LB, Bull FC et al. Global physical activity levels: surveillance progress, pitfalls, and prospects. *The Lancet.* 2012;380(9838):247-57.
4. Lee I-M, Shiroma EJ, Lobelo F et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *The Lancet.* 2012;380(9838):219-29.
5. Ding D, Lawson KD, Kolbe-Alexander TL et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *The Lancet.* 2016;388(10051):1311-24.
6. Das P, Horton R. Physical activity-time to take it seriously and regularly. *The Lancet.* 2016;388(10051):1254.
7. Ekelund U, Steene-Johannessen J, Brown WJ et al. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *The Lancet.* 2016;388(10051):1302-10.
8. Reis RS, Salvo D, Ogilvie D et al. Scaling up physical activity interventions worldwide: stepping up to larger and smarter approaches to get people moving. *The Lancet.* 2016;388(10051):1337-48.
9. Sallis JF, Bull F, Guthold R et al. Progress in physical activity over the Olympic quadrennium. *The Lancet.* 2016;388(10051):1325-36.
10. American College of Sports Medicine. Guidelines for Exercise Testing and Prescription. 10th ed. Philadelphia (PA): Lippincott Williams & Wilkins; 2018: 22-43.
11. American College of Sports Medicine. Guidelines for Exercise Testing and Prescription. 9th ed. Philadelphia (PA): Lippincott Williams & Wilkins; 2014:19-34.
12. Riebe D, Franklin BA, Thompson PD et al. Updating ACSM's recommendations for exercise preparticipation health screening. *Med Sci Sports Exerc.* 2015;47(11):2473-9.
13. Whitfield GP, Riebe D, Magal M, Liguori G. Applying the ACSM Preparticipation Screening Algorithm to US Adults: National Health and Nutrition Examination Survey 2001-2004. *Med Sci Sports Exerc.* 2017;49(10):2056-63.

14. Malina RM. Physical activity and fitness: pathways from childhood to adulthood. *American Journal of Human Biology*. 2001;13(2):162-72.
15. Whitfield GP, Gabriel KKP, Rahbar MH, Kohl HW. Application of the American Heart Association/American College of Sports Medicine Adult Preparticipation Screening Checklist to a nationally representative sample of US adults aged 9=40 years from the National Health and Nutrition Examination Survey 2001 to 2004. *Circulation*. 2014;129(10):1113–20.
16. Rognmo Ø, Moholdt T, Bakken H et al. Cardiovascular risk of high-versus moderate-intensity aerobic exercise in coronary heart disease patients. *Circulation*. 2012; 112.123117.
17. Siscovick DS, Weiss NS, Fletcher RH, Lasky T. The incidence of primary cardiac arrest during vigorous exercise. *N Engl J Med*. 1984;311(14):874-7.
18. Thompson PD, Funk EJ, Carleton RA, Sturner WQ. Incidence of death during jogging in Rhode Island from 1975 through 1980. *JAMA*. 1982;247(18):2535-8.
19. Shephard RJ. Limits to the measurement of habitual physical activity by questionnaires. *British journal of sports medicine*. 2003;37(3):197-206.
20. Prince SA, Adamo KB, Hamel ME, Hardt J, Gorber SC, Tremblay M. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *International Journal of Behavioral Nutrition and Physical Activity*. 2008;5(1):56.
21. Hill K, Gardiner P, Cavalheri V, Jenkins S, Healy G. Physical activity and sedentary behaviour: applying lessons to chronic obstructive pulmonary disease. *Internal medicine journal*. 2015;45(5):474-82.
22. Asthma UK: <https://www.asthma.org.uk/about/media/facts-and-statistics/>. Date accessed 5<sup>th</sup> November 2018.
23. Lötvall J, Akdis CA, Bacharier LB et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *Journal of Allergy and Clinical Immunology*. 2011;127(2):355-60.
24. Price OJ, Hull JH, Ansley L. Advances in the diagnosis of exercise-induced bronchoconstriction. *Expert review of respiratory medicine*. 2014;8(2):209-20.
25. Nici L, Donner C, Wouters E et al. American thoracic society/European respiratory society statement on pulmonary rehabilitation. *American journal of respiratory and critical care medicine*. 2006;173(12):1390-413.

- 391 26. Bolton CE, Bevan-Smith EF, Blakey JD et al. British Thoracic Society guideline on  
392 pulmonary rehabilitation in adults: accredited by NICE. *Thorax*. 2013;68(Suppl 2):ii1-  
393 ii30.
- 394 27. Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global  
395 epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of  
396 prevalence, incidence, and outcomes. *Hepatology*. 2016;64(1):73-84.
- 397 28. Vestbo J, Hurd SS, Agustí AG et al. Global strategy for the diagnosis, management,  
398 and prevention of chronic obstructive pulmonary disease: GOLD executive summary.  
399 *American journal of respiratory and critical care medicine*. 2013;187(4):347-65.
- 400 29. Yano Y, Stamler J, Garside DB et al. Isolated systolic hypertension in young and  
401 middle-aged adults and 31-year risk for cardiovascular mortality: the Chicago Heart  
402 Association Detection Project in Industry study. *Journal of the American College of*  
403 *Cardiology*. 2015;65(4):327-35.
- 404 30. World Health Statistics 2018: [tinyurl.com/y8rwczhh](https://tinyurl.com/y8rwczhh). Date accessed: 5<sup>th</sup> November  
405 2018.
- 406  
407  
408  
409  
410  
411